

Treatment Effect Bounds: An Application to Swan-Ganz Catherization*

Jay Bhattacharya
School of Medicine
Stanford University
jay@stanford.edu

Azeem M. Shaikh
Department of Economics
University of Chicago
amshaikh@uchicago.edu

Edward Vytlacil
Department of Economics
Yale University
edward.vytlacil@yale.edu

February 20, 2008

Abstract

We reanalyze data from the observational study by Connors et al. (1996) on the impact of Swan-Ganz catheterization on mortality outcomes. The Connors et al. (1996) study assumes that there are no unobserved differences between patients who are catheterized and patients who are not catheterized and finds that catheterization increases patient mortality. We instead allow for such differences between patients by implementing both the bounds of Manski (1990), which only exploits an instrumental variable, and the bounds of Shaikh and Vytlacil (2005), which exploit mild nonparametric, structural assumptions in addition to an instrumental variable. We propose and justify the use of indicators of weekday admission as an instrument for catheterization in this context. We find that in our application, the Manski (1990) bounds do not indicate whether catheterization increases or decreases mortality, whereas the Shaikh and Vytlacil (2005) bounds reveal that catheterization increases mortality at 30 days and beyond. We also extend the analysis of Shaikh and Vytlacil (2005) to exploit a further nonparametric, structural assumption – that doctors catheterize individuals with systematically worse latent health – and find that this assumption further narrows these bounds and strengthens our conclusions.

*We thank Hidehiko Ichimura, Seung-Hyun Hong and seminar participants at Brigham Young University, University of Chicago, Harvard/MIT, Michigan Ann-Arbor, Michigan State, the ZEW 2nd Conference on Evaluation Research in Mannheim, and the “Applications of Recent Methodologies in Project Evaluation” session at the Winter 2008 ASSA meeting in New Orleans for helpful comments. This research was supported by NSF Grant SES-05-51089.

1 Introduction

We reanalyze data from a well known observational study by Connors et al. (1996) on the impact of Swan-Ganz catheterization on mortality outcomes. The Swan-Ganz catheter is a device placed in patients in the intensive care unit (ICU) to guide therapy. Connors et al. (1996) examine data on mortality outcomes among a population of patients admitted to the ICU and reach the controversial conclusion that patients who receive Swan-Ganz catheterization during their first day in the ICU are 1.27 times more likely to die within 180 days of their admission. Even at 7 days after ICU admission, Connors et al. (1996) find that catheterization increases mortality. This conclusion was very surprising to ICU doctors, many of whom continue to use the Swan-Ganz catheter to guide therapy in the ICU.

The statistical strategy used by Connors et al. (1996) – the propensity score matching method – assumes away the possibility of unobserved differences between catheterized and non-catheterized patients. Our analysis, by comparison, permits the possibility of unobserved differences. We rely on an instrument for Swan-Ganz catheterization to bound the average effect of catheterization on mortality. We consider the bounds of Shaikh and Vytlačil (2005), which exploit not only the instrumental variable, but also threshold crossing properties for both the treatment and the outcome variables. The assumptions underlying these bounds are therefore stronger than those underlying the bounds of Manski (1990). We also extend the analysis of Shaikh and Vytlačil (2005) to exploit the assumption that doctors are catheterizing those patients who have the worst latent health.

We use the day of the week that the patient was admitted to the ICU as an instrument for Swan-Ganz catheterization. This same variable has been used as an instrument for treatment by Hamilton et al. (2000) in their study of the effect of queuing time on mortality in a Canadian population undergoing hip-fracture surgery. We argue that this variable meets the two crucial requirements for an instrument’s validity. First, it is strongly correlated with the application of the treatment: on weekends, patients are less likely to be catheterized. Second, within observable risk classes, it is uncorrelated with outcomes; that is, mortality rates have little to do with the particular day of the week that a patient is admitted to the ICU and more to do with the arc of the patient’s medical condition.

We find that the bounds of Manski (1990) do not permit us to say whether catheterization increases or decreases mortality—stronger assumptions are needed. In contrast, our application of the bounds of Shaikh and Vytlačil (2005), which imposes mild structural assumptions in addition to those required by Manski (1990), shows that Swan-Ganz catheterization increases

mortality at 30 days after catheterization and beyond. Imposing the additional assumption that doctors catheterize individuals with the worst latent health further narrows these bounds.

2 Background on Swan-Ganz Catheterization

The placement of Swan-Ganz catheters is common among ICU patients – over 2 million patients in North America are catheterized each year. A Swan-Ganz catheter is a slender tube with sensors that measures hemodynamic pressures in the right side of the heart and in the pulmonary artery. Once in place, the catheter is often left in place for days, so it can continuously provide information to ICU doctors. This information is often used to make decisions about treatment, such as whether to give the patient medications that affect the functioning of the heart. While there are some risks associated with the placement of the catheter itself, such complications are rare. Rather, the greater risk may come from successful catheter placement. Information from Swan-Ganz catheterization may, for example, lead to false diagnoses of heart failure, which in turn may lead doctors to administer inappropriate treatments.

Before Connors et al. (1996), Gore et al. (1985) and Zion et al. (1990) also found that catheterization increases mortality. Dalen (2001) criticized both studies because they did not control for clinically important differences between the patients who had catheters placed and those who did not. The Connors et al. (1996) study was conceived in part as a response to this criticism. They included a dizzying array of clinical variables designed to control as exhaustively as possible for observed differences between catheterized and non-catheterized patients. In addition, Connors et al. (1996) expanded the set of ICU patients beyond just heart attack patients to all ICU patients. Ironically, Weil (1998) argued that because Connors et al. (1996) expanded the set of patients considered, they failed to take account of important unobserved clinical variables in their statistical work.

Despite substantial criticism, the publication of the Connors et al. (1996) study was seminal in the Swan-Ganz catheterization literature. Subsequent studies have focused on expanding the set of ICU patients considered in the analysis and on minimizing the possibility of selection bias. There has been one reanalysis of the Connors et al. (1996) study. Hirano and Imbens (2001) modify the propensity score matching method by using a model selection procedure to determine which regressors to include in propensity score model. Their main finding is that the Connors et al. (1996) conclusion that catheterization increases mortality risk is robust to their model selection exercise.

Prior to Connors et al. (1996), attempts to organize a randomized trial failed because doctors refused to recruit patients into the control group. The belief in the efficacy of catheterization was so strong that doctors believed it unethical to deny this procedure to patients on the basis of chance—see Fowler and Cook (2003) and Guyatt (1991). Since Connors et al. (1996), there have been at least two randomized trials on specialized ICU populations: Sandham et al. (2003) and Richard et al. (2003). Neither finds statistically significant differences in mortality between catheterized and non-catheterized patients. While it would be appealing to compare our results with these trials, substantial differences between the populations studied in the trials and this study preclude a direct comparison.

In recent work motivated in part by this paper, Li et al. (2008b) and Altonji et al. (2008) also use the data of Connors et al. (1996) to evaluate the effect of Swan-Ganz catheterization. Li et al. (2008b) apply the methodology of Li et al. (2008a), which, like Connors et al. (1996), rules out the possibility of selection on unobservable characteristics. Altonji et al. (2008) use the methodology of Altonji et al. (2005), which allows for selection on unobservable characteristics, but involves restrictions on the relationship between the strength of selection on observable characteristics and the strength of selection on unobservable characteristics. In contrast to these papers, our methodology allows for selection on unobservable characteristics and requires no restrictions on the strength of selection on these characteristics, but, in order to do so, it requires an instrumental variable.

3 Notation and Assumptions

In this section, we define the notation and assumptions. Let Y be an indicator for patient death within the given number of days after admission into the ICU unit, and let D be an indicator for catheterization. Let X be observed individual characteristics determining mortality and let Z be observed individual characteristics determining catheterization. We assume that both Y and D are determined by threshold crossing models; that is,

$$\begin{aligned} Y^* &= r(X, D) - \epsilon \\ Y &= \mathbf{1}\{Y^* \geq 0\} \end{aligned} \tag{1}$$

$$\begin{aligned} D^* &= s(Z) - \nu \\ D &= \mathbf{1}\{D^* \geq 0\} , \end{aligned} \tag{2}$$

where $\mathbf{1}\{A\}$ is the indicator function of the event A and ϵ and ν are unobserved random variables. The latent index Y^* may be interpreted as an unobserved measure of health status,

and the latent index D^* may be interpreted as an unobserved measure of the desire by hospital staff to conduct the catheterization.

Let Y_1 denote the outcome that would be observed if the individual receives treatment, and let Y_0 denote the outcome that would be observed if the individual does not receive treatment. In our framework, these potential outcomes are given by

$$\begin{aligned} Y_1 &= \mathbf{1}\{r(X, 1) - \epsilon \geq 0\} \\ Y_0 &= \mathbf{1}\{r(X, 0) - \epsilon \geq 0\} . \end{aligned}$$

The effect of catheterization on mortality is $Y_1 - Y_0$, and the average effect of the catheterization on mortality is $E[Y_1 - Y_0] = \Pr\{Y_1 = 1\} - \Pr\{Y_0 = 1\}$. Only Y_1 is observed for individuals who receive catheterization, and only Y_0 is observed for individuals who did not receive catheterization.

We assume further that $(X, Z) \perp\!\!\!\perp (\epsilon, \nu)$. We thus allow catheterization to be endogenous, reflecting the possible dependence between ϵ and ν , but we assume that all other regressors are exogenous. We also assume that (ϵ, ν) has a strictly positive density with respect to Lebesgue measure on \mathbf{R}^2 . This assumption eases the exposition but is not essential. We also require that there is at least one variable in Z that is not in X ; that is, there is some variable that affects the decision to perform catheterization, but does not directly affect mortality. Such a variable is often referred to as an instrumental variable. In our application, we will use an indicator variable for whether the patient was admitted into the ICU on a weekend (rather than a weekday) for this purpose.

Remark 3.1 Vytlacil (2002) establishes the equivalence between the threshold crossing model on D defined in (2) and the monotonicity assumption of Imbens and Angrist (1994). We impose the threshold crossing structure on the equations for both Y and D . Equivalently, we impose the monotonicity assumption of Imbens and Angrist (1994) on both Y and D . ■

Remark 3.2 An important special case of our model is the bivariate probit model with structural shift of Heckman (1978), which imposes the further assumptions that $r(X, D) = X\beta + D\alpha$, $s(Z) = Z\gamma$, and (ϵ, ν) is distributed as a bivariate normal variable with zero means and unit variances. Our model nests this model as a special case, but does not require any of its parametric assumptions. ■

4 Bounds on the Average Treatment Effect

In this section, we develop several different bounds on the average treatment effect. For ease of exposition, suppose that there are no X covariates and that Z is a binary random variable. See Remark 4.5 for a discussion of how the results below would change if these assumptions were relaxed. We assume further that Z is ordered so that $\Pr\{D = 1|Z = 1\} > \Pr\{D = 1|Z = 0\}$. In our application, $Z = 1$ therefore corresponds to a admission into an ICU on a weekday while $Z = 0$ corresponds to admission on a weekend.

4.1 Bounds of Manski (1990)

Manski (1990) only assumes that Y_1 and Y_0 are mean independent of Z ; that is, $\Pr\{Y_0 = 1 | Z\} = \Pr\{Y_0 = 1\}$ and $\Pr\{Y_1 = 1 | Z\} = \Pr\{Y_1 = 1\}$. Note that

$$\Pr\{Y_1 = 1 | Z = z\} = \Pr\{D = 1, Y_1 = 1 | Z = z\} + \Pr\{D = 0, Y_1 = 1 | Z = z\}.$$

Since $Y = Y_1$ when $D = 1$, $\Pr\{D = 1, Y_1 = 1 | Z = z\} = \Pr\{D = 1, Y = 1 | Z = z\}$ is immediately identified from the distribution of the observed data. $\Pr\{D = 0, Y_1 = 1 | Z = z\} = \Pr\{D = 0 | Z = z\}\Pr\{Y_1 = 1 | D = 0, Z = z\}$, on the other hand, is not identified from the distribution of the observed data since we never observe Y_1 for individuals with $D = 0$. But $0 \leq \Pr\{Y_1 = 1 | D = 0, Z = z\} \leq 1$, so

$$\begin{aligned} \Pr\{D = 1, Y = 1|Z = z\} &\leq \Pr\{Y_1 = 1|Z = z\} \\ &\leq \Pr\{D = 1, Y = 1|Z = z\} + \Pr\{D = 0|Z = z\}. \end{aligned}$$

The same argument *mutatis mutandis* can be used to derive similar bounds on $\Pr\{Y_0 = 1|Z = z\}$. Since Y_0 and Y_1 are independent of Z by assumption, we have

$$B_M^L \leq E[Y_1 - Y_0] \leq B_M^U$$

where

$$\begin{aligned} B_M^L &= \max_z \{\Pr\{D = 1, Y = 1|Z = z\}\} \\ &\quad - \min_z \{\Pr\{D = 0, Y = 1|Z = z\} + \Pr\{D = 1|Z = z\}\} \\ B_M^U &= \min_z \{\Pr\{D = 1, Y = 1|Z = z\} + \Pr\{D = 0|Z = z\}\} \\ &\quad - \max_z \{\Pr\{D = 0, Y = 1|Z = z\}\}. \end{aligned}$$

4.2 Bounds of Shaikh and Vytlacil (2005)

Shaikh and Vytlacil (2005) impose the assumptions described in Section 3. Their assumptions, while remaining nonparametric in nature, are stronger than those imposed by Manski (1990).

Under these assumptions,

$$\begin{aligned} \Pr\{Y = 1 \mid Z = z\} &= \Pr\{D = 1, Y = 1 \mid Z = z\} + \Pr\{D = 0, Y = 1 \mid Z = z\} \\ &= \Pr\{D = 1, Y_1 = 1 \mid Z = z\} + \Pr\{D = 0, Y_0 = 1 \mid Z = z\} \\ &= \Pr\{\nu \leq s(z), \epsilon \leq r(1)\} + \Pr\{\nu > s(z), \epsilon \leq r(0)\} . \end{aligned}$$

Recall that we have ordered Z so that $\Pr\{D = 1 \mid Z = 1\} > \Pr\{D = 1 \mid Z = 0\}$, which, under our assumptions, implies $s(1) > s(0)$. Thus, if $r(1) > r(0)$,

$$\Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} = \Pr\{s(0) < \nu \leq s(1), r(0) < \epsilon \leq r(1)\} ,$$

and if $r(1) < r(0)$ then

$$\Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} = -\Pr\{s(0) < \nu \leq s(1), r(1) < \epsilon \leq r(0)\} .$$

It follows that

$$\begin{aligned} \Pr\{Y = 1 \mid Z = 1\} > \Pr\{Y = 1 \mid Z = 0\} &\iff r(1) > r(0) \\ \Pr\{Y = 1 \mid Z = 1\} < \Pr\{Y = 1 \mid Z = 0\} &\iff r(1) < r(0) . \end{aligned}$$

Note that $r(1) \geq r(0)$ implies that $Y_1 \geq Y_0$ and $r(1) \leq r(0)$ implies that $Y_1 \leq Y_0$. It follows that if $\Pr\{Y = 1 \mid Z = 1\} \geq \Pr\{Y = 1 \mid Z = 0\}$, for example, then

$$\begin{aligned} \Pr\{Y = 1 \mid D = 1, Z = z\} &\geq \Pr\{Y_0 = 1 \mid D = 1, Z = z\} \\ \Pr\{Y = 1 \mid D = 0, Z = z\} &\leq \Pr\{Y_1 = 1 \mid D = 0, Z = z\} \end{aligned}$$

The resulting bounds on the average treatment effect are given by

$$B_{SV}^L \leq E[Y_1 - Y_0] \leq B_{SV}^U ,$$

where

$$\begin{aligned} B_{SV}^L &= \Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} \\ B_{SV}^U &= \Pr\{D = 1, Y = 1 \mid Z = 1\} + \Pr\{D = 0 \mid Z = 1\} - \Pr\{D = 0, Y = 1 \mid Z = 0\} \end{aligned}$$

when $\Pr\{Y = 1 \mid Z = 1\} > \Pr\{Y = 1 \mid Z = 0\}$,

$$\begin{aligned} B_{SV}^L &= \Pr\{D = 1, Y = 1 \mid Z = 1\} - \Pr\{D = 0, Y = 1 \mid Z = 0\} - \Pr\{D = 1 \mid Z = 0\} \\ B_{SV}^U &= \Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} \end{aligned}$$

when $\Pr\{Y = 1 \mid Z = 1\} < \Pr\{Y = 1 \mid Z = 0\}$, and $B_{SV}^L = B_{SV}^U = 0$ when $\Pr\{Y = 1 \mid Z = 1\} = \Pr\{Y = 1 \mid Z = 0\}$.

Remark 4.1 The Shaikh and Vytlacil (2005) bounds always lie on one side of zero, unless $\Pr\{Y = 1 \mid Z = 1\} = \Pr\{Y = 1 \mid Z = 0\}$, in which case the average treatment effect is identified to be zero. To see this, note that if $\Pr\{Y = 1 \mid Z = 1\} > \Pr\{Y = 1 \mid Z = 0\}$, then the lower bound on the average treatment effect is $\Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} > 0$. Conversely, if $\Pr\{Y = 1 \mid Z = 1\} < \Pr\{Y = 1 \mid Z = 0\}$, then the upper bound on the average treatment effect is $\Pr\{Y = 1 \mid Z = 1\} - \Pr\{Y = 1 \mid Z = 0\} < 0$. The bounds of Shaikh and Vytlacil (2005) therefore always identify the sign of the average treatment effect. ■

Remark 4.2 Under the assumptions that D is given by (2) and that the unobservables are independent of Z , it follows from the Theorem 2 of Heckman and Vytlacil (2001) that the bounds of Manski (1990) may be written as

$$\begin{aligned} B_M^L &= \Pr\{D = 1, Y = 1 \mid Z = 1\} - \Pr\{D = 0, Y = 1 \mid Z = 0\} - \Pr\{D = 1 \mid Z = 0\} \\ B_M^U &= \Pr\{D = 1, Y = 1 \mid Z = 1\} + \Pr\{D = 0 \mid Z = 1\} - \Pr\{D = 0, Y = 1 \mid Z = 0\} . \end{aligned}$$

Note that if $\Pr\{Y = 1 \mid Z = 1\} \geq \Pr\{Y = 1 \mid Z = 0\}$, then $B_{SV}^U = B_M^U$. The upper bounds on the average treatment effect is therefore the same. On the other hand,

$$\begin{aligned} B_{SV}^L - B_M^L &= \Pr\{D = 0, Y = 1 \mid Z = 1\} - \Pr\{D = 1, Y = 1 \mid Z = 0\} \\ &\quad + \Pr\{D = 1 \mid Z = 0\} \\ &= \Pr\{D = 0, Y = 1 \mid Z = 1\} + \Pr\{D = 1, Y = 0 \mid Z = 0\} \geq 0 , \end{aligned}$$

so $B_{SV}^L \geq B_M^L$. Typically, the inequality will in fact be strict. Conversely, if $\Pr\{Y = 1 \mid Z = 1\} \leq \Pr\{Y = 1 \mid Z = 0\}$, then $B_{SV}^L = B_M^L$ and $B_{SV}^U \leq B_M^U$. The bounds of Shaikh and Vytlacil (2005) are therefore smaller than those of Manski (1990). ■

Remark 4.3 Manski and Pepper (2000) consider a “monotone instrumental variables” (MIV) assumption and a “monotone treatment response” (MTR) assumption. The MIV assumption is a weaker form of the instrumental variable assumption found in Manski (1990). The MTR assumption requires that one knows *a priori* that $Y_1 \geq Y_0$ for all individuals or that one knows *a priori* that $Y_0 \geq Y_1$ for all individuals. In the present context of the effect of catheterization on mortality, where much of the debate focuses on whether the average effect of catheterization is positive, negative, or zero, imposing MTR is unpalatable since it would involve imposing the

answer to the question of interest. The relationship of our analysis with the analysis of Manski and Pepper (2000) is analyzed in Bhattacharya et al. (2008). As discussed by Bhattacharya et al. (2008), it is not possible to determine the sign of the treatment effect in the same way as Shaikh and Vytlacil (2005) under the assumptions of Manski and Pepper (2000). Current work by Machado et al. (2008) develops the sharp bounds for the average treatment effect under the restriction that the outcome is monotone in the treatment, but without assuming the direction of the monotonicity *a priori* or that the treatment is monotone in the instrument. See also Blundell et al. (2007) for related analysis in the context of bounding changes in the distribution of wages when wages are only observed for workers while allowing for non-random selection into employment. ■

4.3 An Extension of Shaikh and Vytlacil (2005)

In this section, we extend the analysis of Shaikh and Vytlacil (2005) to exploit the additional assumption that doctors catheterize individuals with the worst latent health. This restriction is analogous to the Manski and Pepper (2000) “monotone treatment selection” restriction and to the stochastic dominance restriction considered by Blundell et al. (2007). Formally, we assume that ϵ and ν are positive quadrant dependent (PQD); that is,

$$\Pr\{\epsilon \leq t_0 \mid \nu \leq t_1\} \geq \Pr\{\epsilon \leq t_0\} \text{ for all } t_0, t_1 .$$

Positive quadrant dependence is a relatively weak measure of positive dependence between two random variables. See Joe (1997) for a discussion of the relationship between positive quadrant dependence and other concepts of positive dependence. Put differently, this assumption requires that individuals with unobserved characteristics that make them more likely to be catheterized (have a low value of ν) are individuals with unobserved characteristics that make them more likely to suffer mortality (have a low values of ϵ).

The PQD assumption implies that

$$\Pr\{\epsilon \leq t_0 \mid \nu \leq t_1\} \geq \Pr\{\epsilon \leq t_0 \mid \nu > t_1\} \text{ for all } t_0, t_1 .$$

It follows that

$$\begin{aligned} \Pr\{Y = 1 \mid D = 1, Z = z\} &= \Pr\{\epsilon \leq r(1) \mid \nu \leq s(z)\} \\ &\geq \Pr\{\epsilon \leq r(1) \mid \nu > s(z)\} \\ &= \Pr\{Y_1 = 1 \mid D = 0, Z = z\} . \end{aligned}$$

Similarly, we have that

$$\Pr\{Y_0 = 1 | D = 1, Z = z\} \geq \Pr\{Y = 1 | D = 0, Z = z\} .$$

It therefore follows from the analysis of the preceding section that if $\Pr\{Y = 1 | Z = 1\} \geq \Pr\{Y = 1 | Z = 0\}$, then

$$\begin{aligned} \Pr\{Y = 1 | D = 1, Z = z\} &\geq \Pr\{Y_1 = 1 | D = 0, Z = z\} \geq \Pr\{Y = 1 | D = 0, Z = z\} \\ \Pr\{Y = 1 | D = 1, Z = z\} &\geq \Pr\{Y_0 = 1 | D = 1, Z = z\} \geq \Pr\{Y = 1 | D = 0, Z = z\} ; \end{aligned}$$

if, on the other hand, $\Pr\{Y = 1 | Z = 1\} \leq \Pr\{Y = 1 | Z = 0\}$, then

$$\begin{aligned} \min\{\Pr\{Y = 1 | D = 1, Z = z\}, \Pr\{Y = 1 | D = 0, Z = z\}\} &\geq \Pr\{Y_1 = 1 | D = 0, Z = z\} \geq 0 \\ \max\{\Pr\{Y = 1 | D = 1, Z = z\}, \Pr\{Y = 1 | D = 0, Z = z\}\} &\leq \Pr\{Y_0 = 1 | D = 1, Z = z\} \leq 1 . \end{aligned}$$

These results bound $\Pr\{Y_0 = 1\}$ and $\Pr\{Y_1 = 1\}$. If, for example, $\Pr\{Y = 1 | Z = 1\} > \Pr\{Y = 1 | Z = 0\}$, then

$$\begin{aligned} \Pr\{Y_1 = 1\} &= \Pr\{Y_1 = 1 | Z = z\} \\ &= \Pr\{D = 1 | Z = z\} \Pr\{Y_1 = 1 | D = 1, Z = z\} \\ &\quad + \Pr\{D = 0 | Z = z\} \Pr\{Y_1 = 1 | D = 0, Z = z\} \\ &\leq \Pr\{Y = 1 | D = 1, Z = z\} , \end{aligned}$$

which implies that

$$\Pr\{Y_1 = 1\} \leq \min_z \{\Pr\{Y = 1 | D = 1, Z = z\}\} .$$

Using arguments given in Shaikh and Vytlačil (2005), it is possible show that

$$\min_z \{\Pr\{Y = 1 | D = 1, Z = z\}\} = \Pr\{Y = 1 | D = 1, Z = 1\} .$$

The bounds resulting from this line of reasoning are given by

$$B_{PQD}^L \leq E[Y_1 - Y_0] \leq B_{PQD}^U ,$$

where

$$\begin{aligned} B_{PQD}^L &= \Pr\{Y = 1 | Z = 1\} - \Pr\{Y = 1 | Z = 0\} \\ B_{PQD}^U &= \Pr\{Y = 1 | D = 1, Z = 1\} - \Pr\{Y = 1 | D = 0, Z = 0\}, \end{aligned}$$

when $\Pr\{Y = 1 \mid Z = 1\} > \Pr\{Y = 1 \mid Z = 0\}$,

$$\begin{aligned} B_{PQD}^L &= \Pr\{D = 1, Y = 1 \mid Z = 1\} - \Pr\{D = 0, Y = 1 \mid Z = 0\} \\ &\quad - \Pr\{D = 1 \mid Z = 0\} \\ B_{PQD}^U &= \Pr\{D = 1, Y = 1 \mid Z = 1\} + \Pr\{D = 0 \mid Z = 1\} \\ &\quad \times \min\{\Pr\{Y = 1 \mid D = 1, Z = 1\}, \Pr\{Y = 1 \mid D = 0, Z = 1\}\} \\ &\quad - \Pr\{D = 0, Y = 1 \mid Z = 0\} - \Pr\{D = 1 \mid Z = 0\} \\ &\quad \times \max\{\Pr\{Y = 1 \mid D = 1, Z = 0\}, \Pr\{Y = 1 \mid D = 0, Z = 0\}\}, \end{aligned}$$

when $\Pr\{Y = 1 \mid Z = 1\} < \Pr\{Y = 1 \mid Z = 0\}$, and $B_{PQD}^L = B_{PQD}^U = 0$ when $\Pr\{Y = 1 \mid Z = 1\} = \Pr\{Y = 1 \mid Z = 0\}$.

Remark 4.4 The PQD bounds are (weakly) narrower than the SV bounds. To see this, first suppose that $\Pr\{Y = 1 \mid Z = 1\} > \Pr\{Y = 1 \mid Z = 0\}$. In this case, $B_{SV}^L = B_{PQD}^L$, but

$$\begin{aligned} B_{SV}^U - B_{PQD}^U &= \Pr\{D = 0 \mid Z = 1\} \times \Pr\{Y = 0 \mid D = 1, Z = 1\} \\ &\quad + \Pr\{D = 1 \mid Z = 0\} \times \Pr\{Y = 1 \mid D = 0, Z = 0\} \geq 0, \end{aligned}$$

so $B_{SV}^U \geq B_{PQD}^U$. Similarly, if $\Pr\{Y = 1 \mid Z = 1\} < \Pr\{Y = 1 \mid Z = 0\}$, then it is possible to show that $B_{SV}^U = B_{PQD}^U$, but $B_{SV}^L \leq B_{PQD}^L$. Typically, these inequalities will in fact be strict. If $\Pr\{Y = 1 \mid Z = 1\} = \Pr\{Y = 1 \mid Z = 0\}$, then the average treatment effect is identified to be zero and the two sets of bounds coincide. ■

Remark 4.5 Throughout Section 4, we have assumed that there are no X covariates and that Z is binary. Relaxing these assumptions is straightforward. If X is contained in Z , then all of the analysis can simply be carried out conditional on X . If, on the other hand, there exists a component of X that is not contained in Z , then it is possible to further narrow the bounds on the average treatment effect. If there is a continuous component of X that is not contained in Z , then it is possible to obtain point identification. If Z is not binary, then all of the analysis can be carried out with z_1 in place of 1 and z_0 in place of 0, where z_1 maximizes $\Pr\{D = 1 \mid Z = z\}$ and z_0 minimizes $\Pr\{D = 1 \mid Z = z\}$. For further details, see Shaikh and Vytlacil (2005) and Vytlacil and Yildiz (2007). ■

5 Estimation and Inference

In this section, we discuss estimation and inference for each of the bounds described in the preceding section. We also briefly discuss a means of testing for the threshold crossing structure

on the treatment equation. For ease of exposition, we assume again that there are no X covariates. We also assume, as in the preceding section, that Z is ordered so that $\Pr\{D = 1|Z = 1\} > \Pr\{D = 1|Z = 0\}$. Let P denote the distribution of (Y, D, Z) and let $(Y_i, D_i, Z_i), i = 1, \dots, n$ be an i.i.d. sample of random variables with distribution P . We assume throughout that P is such that $0 < \Pr\{D = d, Y = y, Z = z\} < 1$ for all values of $(d, y, z) \in \{0, 1\}^3$.

5.1 Estimation

5.1.1 Bounds of Manski (1990)

Let

$$n_z = |\{1 \leq i \leq n : Z_i = z\}| \quad (3)$$

and define

$$\begin{aligned} \hat{B}_{M,n}^L &= \max_z \left\{ \frac{1}{n_z} \sum_{1 \leq i \leq n: Z_i=z} D_i Y_i \right\} - \min_z \left\{ \frac{1}{n_z} \sum_{1 \leq i \leq n: Z_i=z} ((1 - D_i)Y_i + D_i) \right\} \\ \hat{B}_{M,n}^U &= \min_z \left\{ \frac{1}{n_z} \sum_{1 \leq i \leq n: Z_i=z} (D_i Y_i + (1 - D_i)) \right\} - \max_z \left\{ \frac{1}{n_z} \sum_{1 \leq i \leq n: Z_i=z} (1 - D_i)Y_i \right\}. \end{aligned}$$

Clearly, $\hat{B}_{M,n}^L \xrightarrow{P} B_M^L$ and $\hat{B}_{M,n}^U \xrightarrow{P} B_M^U$, where B_M^L and B_M^U are as defined in Section (4.1). It follows that $[\hat{B}_{M,n}^L, \hat{B}_{M,n}^U]$ converges in probability to $[B_M^L, B_M^U]$ under the Hausdorff metric on subsets of \mathbf{R} .

5.1.2 Bounds of Shaikh and Vytlacil (2005)

Let n_z be given by (3) and define

$$\Delta_n = \left\{ \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} Y_i \right\}. \quad (4)$$

Let $\epsilon_n > 0$ be a sequence of numbers tending to 0, but satisfying $\sqrt{n}\epsilon_n \rightarrow \infty$. Define

$$\begin{aligned} \hat{B}_{SV,n}^L &= \Delta_n \\ \hat{B}_{SV,n}^U &= \left\{ \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} (D_i Y_i + (1 - D_i)) - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} (1 - D_i)Y_i \right\} \end{aligned}$$

when $\Delta_n > \epsilon_n$,

$$\begin{aligned}\hat{B}_{SV,n}^L &= \left\{ \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} D_i Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} ((1-D_i)Y_i + D_i) \right\} \\ \hat{B}_{SV,n}^U &= -\Delta_n\end{aligned}$$

when $\Delta_n < -\epsilon_n$, and $\hat{B}_{SV,n}^L = \hat{B}_{SV,n}^U = 0$ when $|\Delta_n| \leq \epsilon_n$. Clearly, $\hat{B}_{SV,n}^L \xrightarrow{P} B_{SV}^L$ and $\hat{B}_{SV,n}^U \xrightarrow{P} B_{SV}^U$, where B_{SV}^L and B_{SV}^U are as defined in Section 4.2. It follows that $[\hat{B}_{SV,n}^L, \hat{B}_{SV,n}^U]$ converges in probability to $[B_{SV}^L, B_{SV}^U]$ under the Hausdorff metric on subsets of \mathbf{R} .

5.1.3 PQD Bounds

Let n_z be given by (3) and define

$$n_{z,d} = |\{1 \leq i \leq n : Z_i = z, D_i = d\}|. \quad (5)$$

Let Δ_n be given by (4), and let $\epsilon_n > 0$ be a sequence of numbers tending to 0, but satisfying $\sqrt{n}\epsilon_n \rightarrow \infty$. Define

$$\begin{aligned}\hat{B}_{PQD,n}^L &= \Delta_n \\ \hat{B}_{PQD,n}^U &= \left\{ \frac{1}{n_{1,1}} \sum_{1 \leq i \leq n: Z_i=1, D_i=1} Y_i - \frac{1}{n_{0,0}} \sum_{1 \leq i \leq n: Z_i=0, D_i=0} Y_i \right\}\end{aligned}$$

when $\Delta_n > \epsilon_n$,

$$\begin{aligned}\hat{B}_{PQD,n}^L &= \left\{ \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} D_i Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} ((1-D_i)Y_i + D_i) \right\} \\ \hat{B}_{PQD,n}^U &= \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} D_i Y_i + \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} (1-D_i) \\ &\quad \times \min \left\{ \left(\frac{1}{n_{1,1}} \sum_{1 \leq i \leq n: Z_i=1, D_i=1} Y_i \right), \left(\frac{1}{n_{1,0}} \sum_{1 \leq i \leq n: Z_i=1, D_i=0} Y_i \right) \right\} \\ &\quad - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} (1-D_i)Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} D_i \\ &\quad \times \max \left\{ \left(\frac{1}{n_{0,1}} \sum_{1 \leq i \leq n: Z_i=0, D_i=1} Y_i \right), \left(\frac{1}{n_{0,0}} \sum_{1 \leq i \leq n: Z_i=0, D_i=0} Y_i \right) \right\}\end{aligned}$$

when $\Delta_n < -\epsilon_n$, and $\hat{B}_{PQD,n}^L = \hat{B}_{PQD,n}^U = 0$ when $|\Delta_n| \leq \epsilon_n$. Again, $\hat{B}_{PQD,n}^L \xrightarrow{P} B_{PQD}^L$ and $\hat{B}_{PQD,n}^U \xrightarrow{P} B_{PQD}^U$. It follows that $[\hat{B}_{PQD,n}^L, \hat{B}_{PQD,n}^U]$ converges in probability to $[B_{PQD}^L, B_{PQD}^U]$ under the Hausdorff metric on subsets of \mathbf{R} .

Remark 5.1 The above estimators for the Shaikh and Vytlačil (2005) bounds and the PQD bounds have some obvious drawbacks. Both are superefficient when the average treatment effect is in fact equal to zero, which suggests that they would behave poorly in a neighborhood of zero. Moreover, for a given sample of size n , there is no restriction on the level of ϵ_n . As a result, in the next section, we focus instead on constructing confidence sets for the average treatment effect, which do not suffer from such undesirable features. ■

Remark 5.2 Unlike the Manski (1990) bounds, the Shaikh and Vytlačil (2005) bounds and the PQD bounds in general cannot be estimated consistently simply by replacing conditional population means with conditional sample means. This is because these bounds are both discontinuous as a function of $\Pr\{Y = 1|Z = 1\} - \Pr\{Y = 1|Z = 0\}$. In particular, it is not possible to set $\epsilon_n = 0$ and maintain consistency of the estimators. To see this, simply note that when $\Pr\{Y = 1|Z = 1\} - \Pr\{Y = 1|Z = 0\} = 0$, the events $\Delta_n > 0$ and $\Delta_n < 0$ each have probability $1/2$ asymptotically. ■

5.2 Inference

In this section, we construct random sets \mathcal{C}_n such that for each θ between the upper and lower bounds

$$\liminf_{n \rightarrow \infty} \Pr\{\theta \in \mathcal{C}_n\} \geq 1 - \alpha . \quad (6)$$

Such confidence sets have been termed by Romano and Shaikh (2006) as confidence regions for identifiable parameters.

Following Romano and Shaikh (2006), our construction will be based upon test inversion. For each $-1 \leq \theta \leq 1$, we will construct a test of the null hypothesis that θ lies between the upper and lower bounds. The confidence region \mathcal{C}_n will then simply be defined to the set of values for θ for which we fail to reject the corresponding test of the null hypothesis. Concretely, define

$$\mathcal{C}_n = \{-1 \leq \theta \leq 1 : T_n(\theta) \leq \hat{c}_n(\theta, 1 - \alpha)\} , \quad (7)$$

where $T_n(\theta)$ is a test statistic for which large values provide evidence against the null hypothesis and $\hat{c}_n(\theta, 1 - \alpha)$ is an appropriate critical value. The critical value $\hat{c}_n(\theta, 1 - \alpha)$ will be constructed

using subsampling. In order to describe the construction, we require some further notation. Let $b = b_n < n$ be a sequence of positive integers tending to infinity, but satisfying $b_n/n \rightarrow 0$. Index by $i = 1, \dots, N_n = \binom{n}{b}$ the different subsets of $\{1, \dots, n\}$ of size b . Denote by $T_{n,b,i}(\theta)$ the test statistic $T_n(\theta)$ computed using only the i th subset of data of size b . Let $\hat{c}_n(\theta, 1 - \alpha)$ denote the (smallest) $1 - \alpha$ quantile of the distribution

$$L_n(x, \theta) = \frac{1}{N_n} \sum_{1 \leq i \leq N_n} I\{T_{n,b,i}(\theta) \leq x\}. \quad (8)$$

Romano and Shaikh (2006) show that \mathcal{C}_n defined by (7) satisfies the coverage property (6) under weak conditions on the distribution of $T_n(\theta)$ under P . In each of the applications below, it is straightforward to show that these conditions hold using arguments similar to those given in Section 3.2 of Romano and Shaikh (2006).

5.2.1 Bounds of Manski (1990)

Let n_z be given by (3) and let $\bar{z} = (z_1, z_2, z_3, z_4)$. Define

$$\begin{aligned} \delta_{1,n}(z_1, z_2) &= \frac{1}{n_{z_1}} \sum_{1 \leq i \leq n: Z_i = z_1} D_i Y_i - \frac{1}{n_{z_2}} \sum_{1 \leq i \leq n: Z_i = z_2} ((1 - D_i) Y_i + D_i) \\ \delta_{2,n}(z_1, z_2) &= \frac{1}{n_{z_3}} \sum_{1 \leq i \leq n: Z_i = z_1} (D_i Y_i + (1 - D_i)) - \frac{1}{n_{z_4}} \sum_{1 \leq i \leq n: Z_i = z_2} (1 - D_i) Y_i \end{aligned}$$

Denote by $\hat{\sigma}(\{W_i : 1 \leq i \leq n\})$ the usual estimate of the standard deviation of the random variables $W_i, i = 1, \dots, n$. If $z_1 \neq z_2$, then define

$$\begin{aligned} s_{1,n}(z_1, z_2) &= \sqrt{\frac{\gamma_{1,n}(z_1)}{n_{z_1}} + \frac{\gamma_{2,n}(z_2)}{n_{z_2}}} \\ s_{2,n}(z_1, z_2) &= \sqrt{\frac{\gamma_{3,n}(z_1)}{n_{z_1}} + \frac{\gamma_{4,n}(z_2)}{n_{z_2}}}, \end{aligned}$$

where

$$\begin{aligned} \gamma_{1,n}(z_1) &= \hat{\sigma}(\{D_i Y_i : Z_i = z_1, 1 \leq i \leq n\}) \\ \gamma_{2,n}(z_1) &= \hat{\sigma}(\{(1 - D_i) Y_i + D_i : Z_i = z_1, 1 \leq i \leq n\}) \\ \gamma_{3,n}(z_1) &= \hat{\sigma}(\{D_i Y_i + (1 - D_i) : Z_i = z_1, 1 \leq i \leq n\}) \\ \gamma_{4,n}(z_1) &= \hat{\sigma}(\{(1 - D_i) Y_i : Z_i = z_1, 1 \leq i \leq n\}); \end{aligned}$$

if $z_1 = z_2$, then define

$$\begin{aligned} s_{1,n}(z_1, z_2) &= \frac{\hat{\sigma}(\{D_i Y_i - (1 - D_i) Y_i - D_i : Z_i = z_1, 1 \leq i \leq n\})}{\sqrt{n_{z_1}}} \\ s_{2,n}(z_1, z_2) &= \frac{\hat{\sigma}(\{D_i Y_i + (1 - D_i) - (1 - D_i) Y_i : Z_i = z_1, 1 \leq i \leq n\})}{\sqrt{n_{z_1}}}. \end{aligned}$$

For $-1 \leq \theta \leq 1$, define

$$T_n(\theta) = \sum_{\bar{z} \in \{0,1\}^4} \left\{ \left(\frac{\delta_{1,n}(z_1, z_2) - \theta}{s_{1,n}(z_1, z_2)} \right)_+^2 + \left(\frac{\theta - \delta_{2,n}(z_3, z_4)}{s_{2,n}(z_3, z_4)} \right)_+^2 \right\},$$

where $(x)_+ = \max\{x, 0\}$.

Remark 5.3 Imbens and Manski (2004) provide confidence regions with the coverage property (6) for partially identified models where the identified set is an interval whose upper and lower endpoints are means or at least behave like means asymptotically. Although the identified set here is also an interval, the upper and lower endpoints do not have this property, so their analysis is not applicable here. ■

5.2.2 Bounds of Shaikh and Vytlacil (2005)

Let Δ_n be given by (4) and define

$$s_n = \sqrt{\frac{\hat{\sigma}(\{Y_i : Z_i = 1, 1 \leq i \leq n\})}{n_1} + \frac{\hat{\sigma}(\{Y_i : Z_i = 0, 1 \leq i \leq n\})}{n_0}}. \quad (9)$$

For $0 < \theta \leq 1$, define

$$T_n(\theta) = \left(\frac{-\Delta_n}{s_n} \right)_+^2 + \left(\frac{\Delta_n - \theta}{s_n} \right)_+^2 + \left(\frac{\theta - \delta_{2,n}(1, 0)}{s_{2,n}(1, 0)} \right)_+^2;$$

for $-1 \leq \theta < 0$, define

$$T_n(\theta) = \left(\frac{\Delta_n}{s_n} \right)_+^2 + \left(\frac{\theta - \Delta_n}{s_n} \right)_+^2 + \left(\frac{\delta_{1,n}(1, 0) - \theta}{s_{1,n}(1, 0)} \right)_+^2;$$

and for $\theta = 0$, define

$$T_n(\theta) = \left(\frac{\Delta_n}{s_n} \right)_+^2.$$

5.2.3 PQD Bounds

Let $n_{z,d}$ be given by (5), let Δ_n be given by (4), and let s_n be given by (9). Define

$$\begin{aligned}\delta_{3,n} &= \frac{1}{n_{1,1}} \sum_{1 \leq i \leq n: Z_i=1, D_i=1} Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} Y_i \\ \delta_{4,n} &= \frac{1}{n_{1,1}} \sum_{1 \leq i \leq n: Z_i=1, D_i=1} Y_i - \frac{1}{n_{0,0}} \sum_{1 \leq i \leq n: Z_i=0, D_i=0} Y_i \\ \delta_{5,n} &= \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} Y_i - \frac{1}{n_{0,0}} \sum_{1 \leq i \leq n: Z_i=0, D_i=0} Y_i,\end{aligned}$$

and

$$\begin{aligned}s_{3,n} &= \sqrt{\frac{\hat{\sigma}(\{Y_i : Z_i = 1, D_i = 1, 1 \leq i \leq n\})}{n_{1,1}} + \frac{\hat{\sigma}(\{Y_i : Z_i = 0, 1 \leq i \leq n\})}{n_0}} \\ s_{4,n} &= \sqrt{\frac{\hat{\sigma}(\{Y_i : Z_i = 1, D_i = 1, 1 \leq i \leq n\})}{n_{1,1}} + \frac{\hat{\sigma}(\{Y_i : Z_i = 0, D_i = 0, 1 \leq i \leq n\})}{n_{0,0}}} \\ s_{5,n} &= \sqrt{\frac{\hat{\sigma}(\{Y_i : Z_i = 1, 1 \leq i \leq n\})}{n_1} + \frac{\hat{\sigma}(\{Y_i : Z_i = 0, D_i = 0, 1 \leq i \leq n\})}{n_{0,0}}}.\end{aligned}$$

For $0 < \theta \leq 1$, define

$$T_n(\theta) = \left(\frac{-\Delta_n}{s_n}\right)_+^2 + \left(\frac{\Delta_n - \theta}{s_n}\right)_+^2 + \left(\frac{\theta - \delta_{4,n}}{s_{4,n}}\right)_+^2;$$

for $-1 \leq \theta < 0$, define

$$\begin{aligned}T_n(\theta) &= \left(\frac{\Delta_n}{s_n}\right)_+^2 + \left(\frac{\delta_{1,n}(1,0) - \theta}{s_{1,n}(1,0)}\right)_+^2 + \left(\frac{\theta - \delta_{3,n}}{s_{3,n}}\right)_+^2 + \\ &\quad \left(\frac{\theta - \delta_{4,n}}{s_{4,n}}\right)_+^2 + \left(\frac{\theta - \Delta_n}{s_n}\right)_+^2 + \left(\frac{\theta - \delta_{5,n}}{s_{5,n}}\right)_+^2;\end{aligned}$$

and for $\theta = 0$, define

$$T_n(\theta) = \left(\frac{\Delta_n}{s_n}\right)_+^2.$$

5.3 A Test of Threshold Crossing

As discussed in Remark 4.2, Heckman and Vytlačil (2001) show that when D is given by (2) and that the unobservables are independent of Z the bounds of Manski (1990) may be written as

$$\begin{aligned}B_M^L &= \Pr\{D = 1, Y = 1 | Z = 1\} - \Pr\{D = 0, Y = 1 | Z = 0\} - \Pr\{D = 1 | Z = 0\} \\ B_M^U &= \Pr\{D = 1, Y = 1 | Z = 1\} + \Pr\{D = 0 | Z = 1\} - \Pr\{D = 0, Y = 1 | Z = 0\}.\end{aligned}$$

It is therefore possible to test whether D is given by (2) and that the unobservables are independent of Z by comparing these expressions for the bounds of Manski (1990) with those stated in equation (4.1). These two expressions will be the same if and only if

$$\begin{aligned} \Pr\{D = 0, Y = 1|Z = 1\} - \Pr\{D = 1 | Z = 1\} &\geq \\ &\Pr\{D = 0, Y = 1|Z = 0\} - \Pr\{D = 1 | Z = 0\} \\ \Pr\{D = 1, Y = 1|Z = 0\} + \Pr\{D = 0 | Z = 0\} &\geq \\ &\Pr\{D = 1, Y = 1|Z = 1\} + \Pr\{D = 0 | Z = 1\} \\ \Pr\{D = 1, Y = 1|Z = 1\} &\geq \Pr\{D = 1, Y = 1|Z = 0\} \\ \Pr\{D = 0, Y = 1|Z = 0\} &\geq \Pr\{D = 0, Y = 1|Z = 1\} . \end{aligned}$$

We now describe one of several possible ways of testing whether these inequalities hold jointly. Let n_z be given by (3) and define

$$\begin{aligned} \psi_{1,n} &= \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} ((1 - D_i)Y_i - D_i) - \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} ((1 - D_i)Y_i - D_i) \\ \psi_{2,n} &= \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} (D_i Y_i + (1 - D_i)) - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} (D_i Y_i + (1 - D_i)) \\ \psi_{3,n} &= \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} D_i Y_i - \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} D_i Y_i \\ \psi_{4,n} &= \frac{1}{n_1} \sum_{1 \leq i \leq n: Z_i=1} (1 - D_i)Y_i - \frac{1}{n_0} \sum_{1 \leq i \leq n: Z_i=0} (1 - D_i)Y_i . \end{aligned}$$

Consider the test statistic $T_n = \sum_{1 \leq i \leq 4} (\psi_{i,n})_+^2$. Large values of this test statistic provide evidence against the null hypothesis that all four of the above inequalities hold. We may again construct a critical value for this test statistic using subsampling as described in the beginning of Section 5.2. The validity of such an approach can be verified as before using arguments similar to those given in Section 3.2 of Romano and Shaikh (2006). One may, of course, also divide each of the $\psi_{i,n}$ by its standard error, as was done for the test statistics in the previous sections.

6 Data

The Connors et al. (1996) data come from ICUs at five prominent hospitals – Duke University Medical Center, Durham, NC; MetroHealth Medical Center, Cleveland, OH; St. Joseph’s

Hospital, Marshfield, WI; and University of California Medical Center, Los Angeles, CA. The study admitted only severely ill patients with one of nine disease conditions: acute respiratory failure, chronic obstructive pulmonary disease, congestive heart failure, cirrhosis, nontraumatic coma, metastatic colon cancer, late-stage non-small cell lung cancer, and multiorgan system failure with malignancy or sepsis. 59.2% of the sample is over the age of 60. Murphy and Cluff (1990) provide a detailed description of patient recruitment procedures, including a list of exclusion criteria. Connors et al. (1996) count a patient as catheterized if the procedure was performed within 24 hours of entering the ICU.

There are 5,735 patients, all of whom were admitted to or transferred to the ICU within 24 hours of entering the hospital. Connors et al. (1996) collected a large amount of information about each patient via standardized medical chart abstraction methods and interviews with patients and patient surrogates. Tables 1 - 3 compare patients who were catheterized during their first day of admission to the ICU with those who were not catheterized. These tables present the p -value from a test of the hypothesis that the means of the variables are equal for catheterized and non-catheterized patients.

Table 1 compares patients on the basis of demographic variables and the primary diagnosis at admission. Catheterized patients are more likely to be male (by 4.2%), privately insured (by 6.1%), richer (less likely to have an income of less than \$11,000 per year by 5.5%), and have more schooling (0.2 years more on average). Of the patients in our sample, 60.9% of non-catheterized patients had a primary diagnosis of acute respiratory failure, while only 46% of our catheterized patients had the same diagnosis.

Table 2 compares patients on the basis of disease history prior to admission and functional status. Catheterized patients are more likely to have had a diagnosis of cardiac disease (by 1.9%), congestive heart failure (by 2.0%), or acute myocardial infarction (by 1.2%) in their medical history. Catheterized patients also have a 3.3% lower two month predicted survival rate upon admission than non-catheterized patients – clearly catheterized patients are observably more ill than non-catheterized patients.

Table 3 compares catheterized and non-catheterized patients' laboratory values at admission, as well as any secondary diagnoses these patients may have had at admission. Among the laboratory values, all the clinically significant and interpretable differences point toward the conclusion that catheterized patients are observably sicker.

Remark 6.1 Because of the large number of comparisons we are making, it is likely that we will reject several hypotheses falsely. We use the multiple testing procedure of Holm (1979) to

make the comparisons while controlling the familywise error rate – the probability of even one false rejection – at level α . Let $\hat{p}_{(1)} \leq \dots \leq \hat{p}_{(s)}$ denote the ordered values of the p -values and let $H_{(1)}, \dots, H_{(s)}$ denote the corresponding null hypotheses. If $\hat{p}_{(1)} \geq \alpha/s$, then the procedure rejects no null hypotheses; otherwise, it rejects null hypotheses $H_{(1)}, \dots, H_{(r)}$, where r is the largest index such that $\hat{p}_{(i)} \leq \alpha/(s - i + 1)$ for all $i \leq r$. This procedure always rejects at least as many null hypotheses as the Bonferroni procedure, which simply rejects any null hypothesis H_i for which the corresponding $\hat{p}_i \leq \alpha/s$.

At level $\alpha = .05$, we find that patients who are catheterized differ from those who are not catheterized along 32 of 60 possible variables. The results are qualitatively similar at level $\alpha = .01$. Hence, even after accounting for the multiplicity of comparisons, we maintain our earlier conclusion that catheterized patients are significantly different than non-catheterized patients. ■

7 Instrumenting with Admission Day

A direct comparison of outcomes between catheterized and non-catheterized patients is unlikely to yield the causal effects of catheterization. Even if a full set of controls were included in the analysis, the results would be unconvincing. If catheterized and non-catheterized patients differ on so many *observed* dimensions, it is unlikely that they do not differ on *unobserved* dimensions as well. See Altonji et al. (2005) for a formal justification of this argument. In this section, we develop suggestive evidence that day-of-the-week of admission is an appropriate instrument to determine the causal effect of catheterization on patient mortality.

7.1 Admission Day of Week Predicts Catheterization

We first establish that patients who are admitted to the ICU on a Saturday, Sunday, or Monday are substantially less likely to be catheterized on the day of admission than patients admitted on other days of the week. The results remain similar if we exclude Monday from the definition of the weekend. Figure 1 shows catheterization rates by day-of-the-week for four important clinical groups, based on primary diagnosis upon ICU admission. For all four groups, the probability of being catheterized decreases on weekends. A t -test of the difference in probability of catheterization between weekend and weekday rejects equality at 0.05 level for all four groups. However, the same is not true for patients with chronic obstructive pulmonary

disease (COPD), cirrhosis, coma, and lung cancer: there is no statistically significant difference, so we drop patients from these groups from all subsequent analysis.

7.2 Patient Health and Day of Week of Admission

If patients admitted to the ICU on a weekday differed systematically from patients admitted on weekends, then day-of-the-week would be a poor instrument since it would be correlated with unobserved determinants of ICU patient mortality such as health status. We believe that there should be no such correlation, since the health crises that precipitate ICU admissions are unlikely to respect distinctions between weekdays and weekends. We now present suggestive evidence in favor of this view.

Tables 4 - 6 divide patients into two groups on the basis of the instrument: those who were admitted on a weekday and those who were admitted on a weekend. The tables present mean values of each variable on which we make comparisons, along with standard deviations of those variables and the p -value from a test of the hypothesis that the means of the variables are equal.

Table 4 compares patients based on demographic variables on the primary diagnosis. We find no statistically significant differences between patients admitted on a weekend and those admitted on a weekday when testing at the $\alpha = 0.01$ level. Similarly Table 5, which compares patients based on disease history and functional status, shows no statistically significant differences between weekend and weekday patients. Importantly, there is little difference between these groups in predicted 2-month mortality, or in acute physiology score. Finally, Table 6 compares weekend and weekday patients on the basis of laboratory tests at admission and on secondary diagnoses. There are no statistically significant differences between the two groups on these bases.

Remark 7.1 We could apply the multiple testing procedure of Holm (1979) to make the comparisons while controlling the familywise error rate at level α , but it suffices to note that the multiple testing procedure will never reject more null hypotheses than the naive testing procedure above. Therefore, even after accounting for the multiplicity of comparisons, we find that there are no statistically significant differences between the two groups. ■

Remark 7.2 Even though health status at admission does not appear to vary by the day of week of admission to the ICU, the death rates will vary if catheterization rates depend on

day of week of admission and mortality is effected by catheterization. Figure 2 shows mean mortality rates at 7, 30, 60, 90, and 180 days after ICU admission for patients admitted on weekends and weekdays as well as 95% confidence intervals around the means. At 30, 60, 90, and 180 days post-ICU admission, mortality rates are significantly higher at level $\alpha = 0.05$ for patients admitted on weekdays than patients admitted on weekends. ■

7.3 Day of Week, Hospital Staffing, and Outcomes

Although non-specialists sometimes find it surprising, it is well known in the health services literature that medical staffing can have a major effect on treatment decisions, including the decision to catheterize a patient. Rapoport et al. (2000), for example, find that patients admitted to ICUs that staff a full time ICU physician are two-thirds less likely to be catheterized than those admitted to ICUs with no full time physician. Whether this fact threatens the validity of our instrument depends upon whether there are unobserved differences in treatment between weekday and weekend admissions, unassociated with catheterization, that help determine patient mortality. If so, then admission day would not be a valid instrument.

Evaluating the importance of differences in treatment between weekend and weekday admissions is complicated by the fact that Swan-Ganz catheterization itself is a gateway to a large number of other treatments. For example, ICU physicians often use the information from catheterization to titrate the dose of inotropic drugs, such as dopamine and dobutamine, which are designed to improve cardiac contractility. These drugs have a narrow therapeutic range, and thus getting the dose right can be the difference between killing or inadequately treating a patient. Since catheterization is less likely on weekends, it would be unsurprising to find decreased use of inotropes on weekends as well. We can accommodate such differences in treatment between weekend and weekday admissions by simply reinterpreting the treatment as catheterization *and all the other treatments it enables or encourages* on mortality, rather than catheterization by itself.

It is possible that weekend-weekday staffing differences, for reasons having nothing to do with catheterization or its downstream consequences, may lead to higher patient mortality. If so, then our instrument would be invalid. Since staffing tends to be sparser on weekends, one would expect that mortality rates would be higher then. In fact, in our data mortality rates are higher on weekdays, which is inconsistent with a direct mortality effect of staffing. Furthermore, several studies have found no evidence that staffing differences explain weekend-weekday mortality differences in ICUs—see Ensminger et al. (2004), Wunsch et al. (2004), and

Dobkin (2003).

8 Results

In this section, we first follow the traditional approach of estimating parametric models that allow us to identify the average treatment effect. In particular, we consider linear models and bivariate probit models. We then report results from the three different bounds described in Section 4.

We analyze outcomes t days after admission to the ICU separately for different values of $t = 7, 30, 60, 90,$ and 180 days. For this reason, we write the outcome Y and the potential outcomes Y_0 and Y_1 as functions of t throughout the remainder of the paper.

8.1 Parametric Models

The panel on the righthand side of Figure 3 shows estimates of the coefficient on D and associated 95% confidence intervals obtained from ordinary least squares (OLS) regression of $Y(t)$ on a constant and D for the different values of t . The panel on the lefthand side shows estimates of the coefficient on D and associated 95% confidence intervals obtained from OLS regression of $Y(t)$ on a constant, D and the full set of covariates listed in Tables 1 - 3 for the different values of t . For these estimates to be consistent for the average treatment effect, we require that the treatment D be independent of the potential outcomes $Y_0(t)$ and $Y_1(t)$ (conditional on covariates) and that the linear model underlying the regression is correct. Under these assumptions, one might conclude from these results that excess mortality due to catheterization 7 days after ICU admission is about two percentage points and it increases to about six percentage points 30 days after admission.

Even if these assumptions are correct, a possible problem with these results is that they ignore the possibility that catheterized patients differ in unobserved ways from non-catheterized patients. For this reason, we may consider maximum likelihood estimation of the bivariate probit model described in Remark 3.2. The bivariate probit model is a common approach in the context of a dummy endogenous regressor as a determinant of a binary outcome; see, for example, Goldman et al. (2001). For the sake of brevity, we report only the results without X regressors.

Figure 4 shows 95% confidence intervals around the average treatment effect $E[Y_1(t) - Y_0(t)]$ from maximum likelihood estimation of the bivariate probit model for the different values of t . Like the OLS results, the bivariate probit results suggest that catheterization increases mortality rates at 30, 60, 90, and 180 days after ICU admission. Like the OLS results without covariates but unlike the OLS results with covariates, the confidence interval for the average treatment effect crosses zero at 7 days.

Although the bivariate probit model allows catheterized and non-catheterized patients to differ in unobserved ways, it still requires many strong, parametric assumptions. We therefore consider the three nonparametric bounds described in Section 4

8.2 Nonparametric Bounds

We consider each of the three bounds described in Section 4. Each of these bounds rely upon our instrumental variable – an indicator for whether the patient was admitted to the ICU on a Tuesday - Friday. For each of the bounds, we display 95% confidence intervals for the average treatment effect computed as described in Section 5. Computational details are described below in Remark 8.4. The lefthand side of Figure 5 shows the bounds of Manski (1990) for the whole sample, whereas the righthand side shows the bounds of Shaikh and Vytlacil (2005) for the whole sample. The Manski (1990) bounds have a width of nearly one and thus always fail to exclude zero; see Remark 8.1 below for further discussion. The Shaikh and Vytlacil (2005) bounds, by contrast, are considerably more informative: at 7 days after ICU admission, these bounds include zero; but at 30 days after admission to the ICU and beyond, the bounds suggest that catheterization increases mortality. The width of the these bounds are about half the width of the Manski (1990) bounds.

Recall that Connors et al. (1996) found that catheterization increases mortality even at 7 days using this same data set that we use here, but a different statistical method that assumes that there are no unobserved differences between catheterized and non-catheterized patients. Their result raises the question of why ICU doctors do not observe the increased mortality from catheterization and react accordingly. The Shaikh and Vytlacil (2005) bounds provides a possible answer – ICU doctors do not see rise in mortality which happens only after many patients have been released from the ICU.

Figure 6 shows the bounds from the extension of Shaikh and Vytlacil (2005) described in Section 4.3. These bounds impose the restriction that doctors are effective at triaging patients so that it is those patients with the worst health who are actually catheterized. These

figures show that imposing this plausible restriction decreases the width of the treatment effect bounds, often dramatically.

By construction, these bounds are always on the same side of zero as the Shaikh and Vytlacil (2005) bounds. The reduction in the width of the bounds is greatest when the average treatment effect is positive; that is, when catheterization increases mortality. This is to be expected, as the PQD restriction rules out the possibility that doctors cause great harm to large numbers of their patients. On the other hand, the PQD bounds have a lower upper bound than the Shaikh and Vytlacil (2005) bounds when the average treatment effect is negative, that is, when catheterization decreases mortality, so it may permit researchers to conclude, for example, that the intervention is cost-effective even as the Shaikh and Vytlacil (2005) bounds permit the possibility that it may not be.

Remark 8.1 Despite the evidence presented in Section 7, it is interesting to consider how our inferences would change if we did not rely upon our instrumental variable. One possible answer is to rely on the bounds of Manski (1990), which may be constructed without an instrument. In that case, the width of the bounds is always exactly one and thus always fail to exclude zero. A second possible answer is given by the analysis of Section 4.3, which may also be constructed without an instrument. In that case, the PQD assumption reduces to $\Pr\{Y_1 = 1 \mid D = 1\} \geq \Pr\{Y_1 = 1 \mid D = 0\}$ and $\Pr\{Y_0 = 1 \mid D = 1\} \geq \Pr\{Y_0 = 1 \mid D = 0\}$, which implies the following bounds on the average treatment effect:

$$\begin{aligned} & \Pr\{Y_1 = 1 \mid D = 1\} \Pr\{D = 1\} - \Pr\{Y_0 = 1 \mid D = 0\} \Pr\{D = 0\} - \Pr\{D = 1\} \\ & \leq E[Y_1 - Y_0] \leq \Pr\{Y_1 = 1 \mid D = 1\} - \Pr\{Y_0 = 1 \mid D = 0\} \end{aligned}$$

Figure 7 shows these bounds and associated 95% confidence intervals. In every case, the bounds cross zero, though their width is substantially less than one. The PQD assumption by itself is therefore not enough to identify the direction of the treatment effect. ■

Remark 8.2 Heckman and Vytlacil (2001) show that the threshold crossing structure implies that $B_M^U - B_M^L = 1 - \Pr\{D = 1 \mid Z = 1\} + \Pr\{D = 1 \mid Z = 0\}$, where Z is ordered such that $\Pr\{D = 1 \mid Z = 1\} > \Pr\{D = 1 \mid Z = 0\}$. If $\Pr\{D = 1 \mid Z = 1\}$ is close to one and $\Pr\{D = 1 \mid Z = 0\}$ is close to zero, then the bounds will have width close to zero. In contrast, if $\Pr\{D = 1 \mid Z = 1\}$ is close to $\Pr\{D = 1 \mid Z = 0\}$, the width of the bounds will be nearly one, i.e., almost as wide as the naive bounds that do not impose or exploit an instrument described in Remark 8.1. Our empirical result that the width of the bounds is close to one

is a direct result of the instrument being weak in the sense that $\Pr\{D = 1|Z = 1\}$ is close to $\Pr\{D = 1|Z = 0\}$. This is a separate issue from the question of whether the instrument is highly statistically significant in the propensity score model. As we discuss in Section 7.1, for the patient groups we analyze, we can reject that $\Pr\{D = 1|Z = 1\} = \Pr\{D = 1|Z = 0\}$. Furthermore, in the bivariate probit model reported above, a test of the significance of the instruments in the catheterization equation rejects that the coefficient on the instruments are jointly zero at $p = 0.0072$ for the model where the main outcome is 30 day mortality. ■

Remark 8.3 We also implement the test of the threshold crossing assumption that is described in Section 5.3. At the $\alpha = 0.10$ level and for each value of t we fail to reject the inequalities shown in that section, providing evidence in favor of the assumptions underlying the bounds described in Sections 4.2 and 4.3. ■

Remark 8.4 For the results we reported above, we used a subsample size of $b = 50$. In results not reported here, we also tried different subsample sizes ranging from 25 to 75 and found that our results are remained similar for these values of b . Finally, because N_n is large, we used an approximation to (8) in which we randomly chose with replacement $B_n = 1000$ of the N_n possible subsamples. It follows from Corollary 2.4.1 of Politis et al. (1999) that critical values constructed in this way remain valid provided that B_n tends to infinity. ■

9 Conclusion

While direct comparisons of the mortality of catheterized and non-catheterized patients lead to the conclusion that catheterization increases mortality, we show evidence that this result is due to profound differences between the catheterized and non-catheterized patients; the former are much more severely ill than the latter.

We provide suggestive evidence that weekday admission can serve as an instrumental variable for catheterization. Patients admitted on a weekday are about four to eight percentage points more likely to be catheterized than patients admitted on a weekend. Yet, weekday and weekend patients appear similar in health status along a large number of dimensions. Exploiting an instrumental variable permits us to address the unobserved differences between catheterized and non-catheterized ICU patients.

We turn to bounding approaches that exploit access to our instrument, including the recent approach introduced by Shaikh and Vytlacil (2005), which we compare with the approach of

Manski (1990). We find that, while the Manski (1990) bounds always straddle zero, the Shaikh and Vytlacil (2005) bounds typically produces a clearer answer – catheterization increases mortality at 30 days and beyond, while at 7 days the average treatment effect may be zero. We extend the analysis of Shaikh and Vytlacil (2005) to exploit a further nonparametric structural assumption – that doctors catheterize individuals with systematically worse latent health – and find that this assumption further narrows these bounds and strengthens these conclusions.

The main theme of the paper is the trade-off induced by the acceptance of potentially unverifiable structural assumptions. If one is willing to accept very strong structural assumptions, such as those underlying the bivariate probit model, one obtains point identification. At the other extreme, if the only structural assumption one accepts is that probabilities lie between zero and one (such as in the Manski (1990) bounds without an instrument), then the width of the bounds on the average treatment effect is exactly one, so it is not possible to determine the sign of the average treatment effect. In between these two extremes, one may accept different nonparametric, structural assumptions, such as the validity of an instrument or threshold crossing models on the outcome or treatment variables, which may not lead to point identification, but may reduce the width of the bounds considerably, as in our empirical example, and are more palatable than the very strong parametric assumptions required for the bivariate probit model.

Our primary substantive finding is that catheterization improves mortality outcomes only in the short run, if at all, and increases it in the long run. This finding is intuitively appealing because it suggests a possible explanation for the fact that many ICU doctors are committed to the use of the Swan-Ganz catheter. Since most ICU patients leave the ICU well before 30 days after admission have elapsed, ICU doctors may never observe the increase in mortality. Our results also suggest a second (not mutually exclusive) possibility: a simple selection story. Catheterization saves the lives, in the short run, of the most severely ill patients, but the deaths of these patient cannot be staved off for long. Disentangling these possibilities will require even more detailed data and further research.

References

Altonji, A., Elder, T., and Taber, C. (2008). Using selection on observed variables to assess bias from unobservables when evaluating swan-ganz catheterization. *American Economic*

- Review, Papers and Proceedings.* forthcoming.
- Altonji, J., Elder, T., and Taber, C. (2005). Selection on observed and unobserved variables: Assessing the effectiveness of catholic schools. *Journal of Political Economy.* forthcoming.
- Bhattacharya, J., Shaikh, A., and Vytlacil, E. (2008). Treatment effect bounds under monotonicity assumptions: An application to swan ganz catheterization. *American Economic Review, Papers and Proceedings.* forthcoming.
- Blundell, R., Gosling, A., Ichimura, H., and Meghir, C. (2007). Changes in the distribution of male and female wages accounting for employment composition using bounds. *Econometrica.*
- Connors, A., Speroff, T., Dawson, N., Thomas, C., Harrell, F., Wagner, D., Desbiens, N., Goldman, L., Wu, A., Califf, R., Fulkerson, W. J., Vidaillet, H., Broste, S. Bellamy, P., Lynn, J., and Knaus, W. (1996). The effectiveness of right heart catheterization in the initial care of critically ill patients. support investigators. *Journal of the American Medical Association*, 276(11):889–897.
- Dalen, J. (2001). The pulmonary arter catheter-friend, foe, or accomplice? *Journal of the American Medical Association*, 286(3):348–350.
- Dobkin, C. (2003). Hospital staffing and inpatient mortality. mimeo, UC Santa Cruz.
- Ensminger, S., Morales, I., Peters, S., Keegan, M., Finkielman, J., Lymp, J., and Afessa, B. (2004). The hospital mortality of patients admitted to the icu on weekends. *Chest*, 126(4):1292–1298.
- Fowler, R. and Cook, D. (2003). The arc of the pulmonary artery catheter. *Journal of the American Medical Association*, 290(20):2732–2734.
- Goldman, D., Bhattacharya, J., Mccaffrey, D., Duan, N., Leibowitz, A., Joyce, G., and Morton, S. (2001). Effect of Insurance on Mortality in an HIV-Positive Population in Care. *Journal of the American Statistical Association*, 96(455).
- Gore, J., Goldberg, R., Spodick, D., Aplert, J., and Dalen, J. (1985). A community-wide assessment of the use of pulmonary artery catheters in patients with acute myocardial infarction: A prospective autopsy study. *Chest*, 88:567–572.
- Guyatt, G. (1991). A randomized trial of right heart catheterization in critically ill patients. *Journal of Intensive Care Medicine*, 6:91–95. for the Ontario Intensive Care Study Group.

- Hamilton, B., Ho, V., and Goldman, D. (2000). Queing for surgery: Is the u.s. or canada worse off? *Review of Economics and Statistics*, 82(2):297–308.
- Heckman, J. (1978). Dummy endogenous variables in a simultaneous equation system. *Econometrica*, 46:931–959.
- Heckman, J. and Vytlacil, E. (2001). Instrumental variables, selection models, and tight bounds on the average treatment effect. In Lechner, M. and Pfeiffer, F., editors, *Econometric Evaluations of Active Labor Market Policies in Europe*. Physica-Verlag, Heidelberg; New York:.
- Hirano, K. and Imbens, G. (2001). Estimation of causal effects using propensity score weighting: An application to data on right heart catheterization. *Health Services and Outcomes Research Methodology*, 2:259–278.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(65-70).
- Imbens, G. and Angrist, J. (1994). Identification and estimation of local average treatment effects. *Econometrica*, 62(467-476).
- Imbens, G. and Manski, C. (2004). Confidence intervals for partially identified parameters. *Econometrica*, 72(1845-1857).
- Joe, H. (1997). *Multivariate Models and Dependence Concepts*. Chapman and Hall, New York.
- Li, Q., Racine, J., and Wooldridge, J. (2008a). Efficient estimation of average treatment effects with mixed categorical and continuous data. *Journal of Business and Economic Statistics*. forthcoming.
- Li, Q., Racine, J., and Wooldridge, J. (2008b). Estimating average treatment effects with continuous and discrete covariates: The case of swan-ganz catheterization. *American Economic Review, Papers and Proceedings*. forthcoming.
- Machado, C., Shaikh, A., and Vytlacil, E. (2008). Instrumental variables, alternative monotonicity restrictions, and identifying the sign of the average treatment effect. unpublished working paper.
- Manski, C. (1990). Nonparametric bounds on treatment effects. *American Economic Review, Papers and Proceedings*, 80:319–323.

- Manski, C. and Pepper, J. (2000). Monotone instrumental variables: With an application to the returns to schooling. *Econometrica*, 68:997–1010.
- Murphy, D. and Cluff, L. (1990). Support: Study to understand prognoses and preferences for outcomes and risks of treatments-study design. *Journal of Clinical Epidemiology*, 43(suppl):1S–123S.
- Politis, D. N., Romano, J. P., and Wolf, M. (1999). *Subsampling*. Springer, New York.
- Rapoport, J., Teres, D., Steingrub, J., Higgins, T., McGee, W., and Lemeshow, S. (2000). Patient characteristics and ICU organizational factors that influence frequency of pulmonary artery catheterization. *Journal of the American Medical Association*.
- Richard, C., Warszawski, J., Anguel, N., Deye, N., Combes, A., Barnoud, D., Boulain, T., Lefort, Y., Fartoukh, M., Baud, F., Boyer, A., Brochard, L., and Teboul, J. (2003). Early use of the pulmonary artery catheter and outcomes in patients with shock and acute respiratory distress syndrome: A randomized controlled trial. *Journal of the American Medical Association*. for the French Pulmonary Artery Catheter Study Group.
- Romano, J. and Shaikh, A. (2006). Inference for identifiable parameters in partially identified econometric models. mimeo, Stanford University and University of Chicago.
- Sandham, J., Hull, R., Brant, R., Knox, L., Pineo, G., Doig, C., Laporta, D., Viner, S., Passerini, L., Devitt, H., Kirby, A., and Jacka, M. (2003). A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients. *New England Journal of Medicine*. for the Canadian Critical Care Clinical Trials Group.
- Shaikh, A. and Vytlacil, E. (2005). Threshold crossing models and bounds on treatment effects: A nonparametric analysis. NBER Technical Working Paper No. 307.
- Vytlacil, E. (2002). Independence, monotonicity, and latent index models: An equivalence result. *Econometrica*, 70(1):331–341.
- Vytlacil, E. and Yildiz, N. (2007). Dummy endogenous variables in weakly separable models. *Econometrica*, 75(3):775–779.
- Weil, M. (1998). The assault on the Swan-Ganz catheter: A case history of constrained technology, constrained bedside clinicians, and constrained monetary expenditures. *Chest*, 113(5):1379–1386.

Wunsch, H., Mapstone, J., Brady, T., Hanks, R., and Rowan, K. (2004). Hospital mortality associated with day and time of admission to intensive care units. *Intensive Care Medicine*, 30(5):895–901.

Zion, M., Balkin, J., Rosenmann, D., Goldbourt, U., Reicher-Reiss, H., Kaplinsky, E., and Behar, S. (1990). Use of pulmonary artery catheters in patients with acute myocardial infarction. analysis of experience in 5,841 patients in the sprint registry. *Chest*, 98:1331–1335.

Table 1: Catheterized vs. Not Catheterized; Demographic and Diagnostic Comparisons

Variable	Not Catheterized	Catheterized	<i>p</i> -value
Age	61.2 [18.1]	60.4 [15.8]	0.1584
Male	54.3(%) [0.498]	58.5(%) [0.493]	0.0052
Black	17.1(%) [0.376]	15.1(%) [0.358]	0.0714
Other Race	6.1(%) [0.240]	6.7(%) [0.251]	0.4066
Years of Education	11.7 [3.1]	11.9 [3.2]	0.0264
No Insurance	5.2(%) [0.222]	6.2(%) [0.242]	0.1371
Private Insurance	28.1(%) [0.449]	34.2(%) [0.474]	< 0.0001
Medicare	26.5(%) [0.442]	23.2(%) [0.422]	0.0094
Medicaid	12.1(%) [0.326]	8.4(%) [0.277]	< 0.0001
Private Insurance & Medicare	21.1(%) [0.408]	22.4(%) [0.417]	0.2730
Family Income < \$11K per year	57.1(%) [0.495]	51.6(%) [0.500]	0.0002
Family Income \$11K-\$25K	20.2(%) [0.401]	20.9(%) [0.407]	0.5526
Family Income \$25K-\$50K	15.1(%) [0.358]	18.6(%) [0.389]	0.0017
Weight (kg)	65.6 [27.9]	72.3 [27.9]	< 0.0001
Dx: Acute Respiratory Failure	60.9(%) [0.488]	46.0(%) [0.499]	< 0.0001
Dx: Congestive Heart Failure	9.5(%) [0.293]	10.6(%) [0.308]	0.2351
Dx: MOSF with malignancy	9.3(%) [0.290]	8.0(%) [0.271]	0.1265
Dx: MOSF with sepsis	20.3(%) [0.402]	35.4(%) [0.478]	< 0.0001
N	2,596 (57%)	1,976 (43%)	

Table 2: Catheterized vs. Not Catheterized; Disease History and Functional Status

Variable	Not Catheterized	Catheterized	<i>p</i> -value
Hx: Cardiac Disease	18.5(%) [0.388]	20.4(%) [0.403]	0.0973
Hx: Congestive Heart Failure	17.8(%) [0.382]	19.8(%) [0.399]	0.0737
Hx: Dementia	12.4(%) [0.329]	7.0(%) [0.255]	< 0.0001
Hx: Psychiatric Condition	8.5(%) [0.279]	4.7(%) [0.211]	< 0.0001
Hx: Chronic Pulmonary Disease	13.7(%) [0.344]	12.1(%) [0.326]	0.1072
Hx: Renal Disease	5.1(%) [0.220]	5.0(%) [0.217]	0.8479
Hx: Liver Disease	3.8(%) [0.192]	4.6(%) [0.209]	0.2125
Hx: GI Bleed	1.6(%) [0.125]	1.6(%) [0.126]	0.9147
Hx: Malignant Cancer	26.9(%) [0.444]	20.9(%) [0.407]	< 0.0001
Hx: Immunological Disease	27.6(%) [0.447]	29.3(%) [0.455]	0.2144
Transferred from another hospital	9.5(%) [0.293]	14.9(%) [0.356]	< 0.0001
Hx: Acute Myocardial Infarction	3.1(%) [0.173]	4.3(%) [0.202]	0.0352
Hx: Non-metastatic Cancer	20.2(%) [0.402]	15.9(%) [0.366]	0.0002
2 Month Predicted Survival	61.6(%) [0.181]	58.3(%) [0.187]	< 0.0001
Duke Activity Scale Index	20.4 [5.5]	20.7 [5.1]	0.0597
Acute Physiology Score	53.2 [18.5]	60.6 [20.1]	< 0.0001
Glasgow Coma Score	17.8 [26.4]	16.1 [24.9]	0.0206
Diastolic Blood Pressure	81.6 [37.6]	68.2 [33.4]	< 0.0001
Do Not Resuscitate Order	13.0(%) 32 [0.336]	6.7(%) [0.251]	< 0.0001

Note: Each entry shows the mean and standard deviation (in brackets) for each variable.

Table 3: Catheterized vs. Not Catheterized; Lab Values and Secondary Diagnoses

Variable	Not Catheterized	Catheterized	<i>p</i> -value
WBC Count	15.6 [12.4]	16.1 [12.1]	0.1895
Heart Rate	115.2 [40.2]	120.4 [39.9]	< 0.0001
Respiratory Rate	30.4 [13.6]	27.0 [14.0]	< 0.0001
Temperature (°C)	37.8 [1.8]	37.7 [1.8]	0.0120
$PAO_2/(0.01*FiO_2)$	235.4 [116.4]	189.6 [104.3]	< 0.0001
Albumin	3.1 [0.7]	3.0 [0.9]	< 0.0001
Hematocrit	31.6 [8.2]	30.3 [7.1]	< 0.0001
Bilirubin	1.9 [4.3]	2.6 [5.2]	< 0.0001
Creatinine	2.1 [2.2]	2.5 [2.1]	< 0.0001
Sodium	137.0 [8.1]	136.3 [7.6]	0.0044
Potassium	4.1 [1.1]	4.0 [1.0]	0.2147
$PACO_2$	38.0 [11.2]	36.4 [10.0]	< 0.0001
Serum Ph	7.4 [0.1]	7.4 [0.1]	0.0002
2nd Dx: Respiratory Dx	42.3(%) [0.494]	28.5(%) [0.451]	< 0.0001
2nd Dx: Orthopedic Dx	0.1(%) [0.028]	0.2(%) [0.045]	0.2461
2nd Dx: Neurological Dx	12.4(%) [0.329]	4.6(%) [0.210]	< 0.0001
2nd Dx: GI Dx	13.1(%) [0.337]	18.6(%) [0.389]	< 0.0001
2nd Dx: Renal Dx	5.0(%) [0.217]	7.0(%) [0.255]	0.0040
2nd Dx: Metabolic Dx	5.7(%) [0.232]	4.2(%) [0.201]	0.0217
2nd Dx: Hematological Dx	8.6(%) [0.280]	5.4(%) [0.226]	< 0.0001
2nd Dx: Sepsis	18.6(%) [0.389]	24.8(%) [0.432]	< 0.0001
2nd Dx: Trauma	0.7(%) [0.081]	1.7(%) [0.128]	0.0011

Note: Each entry shows the mean and standard deviation (in brackets) for each variable.

Table 4: Weekend vs. Weekday Admissions; Demographics and Diagnoses

Variable	Weekday	Weekend	<i>p</i> -value
Age	60.7 [17.2]	61.1 [17.1]	0.4028
Male	55.9(%) [0.497]	56.4(%) [0.496]	0.7526
Black	15.7(%) [0.364]	16.9(%) [0.375]	0.2730
Other Race	6.4(%) [0.244]	6.4(%) [0.245]	0.9630
Years of Education	11.7 [3.1]	11.8 [3.2]	0.1297
No Insurance	6.1(%) [0.239]	5.0(%) [0.218]	0.1275
Private Insurance	30.3(%) [0.460]	31.3(%) [0.464]	0.4804
Medicare	25.0(%) [0.433]	25.3(%) [0.435]	0.8292
Medicaid	10.7(%) [0.310]	10.1(%) [0.302]	0.5148
Private Insurance & Medicare	21.5(%) [0.411]	21.8(%) [0.413]	0.8023
Family Income < \$11K per year	55.9(%) [0.497]	53.0(%) [0.499]	0.0533
Family Income \$11K-\$25K	20.2(%) [0.402]	20.9(%) [0.407]	0.5442
Family Income \$25K-\$50K	16.1(%) [0.367]	17.5(%) [0.380]	0.2185
Weight (kg)	68.8 [28.4]	68.0 [27.6]	0.3994
Dx: Acute Respiratory Failure	53.4(%) [0.499]	56.1(%) [0.496]	0.0707
Dx: Congestive Heart Failure	10.8(%) [0.311]	8.8(%) [0.283]	0.0221
Dx: MOSF with malignancy	9.0(%) [0.286]	8.4(%) [0.277]	0.4915
Dx: MOSF with sepsis	26.9(%) [0.443]	26.8(%) [0.443]	0.9642
N	2,699 (59%)	1,873 (41%)	

Table 5: Weekend vs. Weekday Admissions; Disease History and Functional Status

Variable	Weekday	Weekend	<i>p</i> -value
Hx: Cardiac Disease	20.8(%) [0.406]	17.2(%) [0.377]	0.0022
Hx: Congestive Heart Failure	20.0(%) [0.400]	16.7(%) [0.373]	0.0049
Hx: Dementia	10.7(%) [0.309]	9.1(%) [0.287]	0.0711
Hx: Psychiatric Condition	7.1(%) [0.257]	6.4(%) [0.245]	0.3513
Hx: Chronic Pulmonary Disease	12.5(%) [0.331]	13.8(%) [0.345]	0.2029
Hx: Renal Disease	4.9(%) [0.215]	5.3(%) [0.224]	0.5112
Hx: Liver Disease	4.6(%) [0.209]	3.5(%) [0.183]	0.0605
Hx: GI Bleed	1.7(%) [0.129]	1.4(%) [0.119]	0.4858
Hx: Malignant Cancer	24.0(%) [0.427]	24.8(%) [0.432]	0.5078
Hx: Immunological Disease	28.4(%) [0.451]	28.2(%) [0.450]	0.8880
Transferred from another hospital	11.6(%) [0.320]	12.2(%) [0.327]	0.5277
Hx: Acute Myocardial Infarction	3.9(%) [0.193]	3.2(%) [0.176]	0.2453
Hx: Non-metatstatic Cancer	18.3(%) [0.387]	18.4(%) [0.388]	0.9203
2 Month Predicted Survival	60.5(%) [0.183]	59.7(%) [0.185]	0.1333
Duke Activity Scale Index	20.4 [5.3]	20.8 [5.5]	0.0144
Acute Physiology Score	56.4 [19.6]	56.4 [19.5]	0.9938
Glasgow Coma Score	17.1 [26.1]	17.0 [25.3]	0.8641
Diastolic Blood Pressure	76.1 [36.3]	75.4 [36.6]	0.5686
Do Not Resuscitate Order	10.6(%) [0.308]	9.8(%) [0.298]	0.3976

35

Note: Each entry shows the mean and standard deviation (in brackets) for each variable.

Table 6: Weekend vs. Weekday Admissions; Lab Values and Secondary Diagnoses

Variable	Weekday	Weekend	<i>p</i> -value
WBC Count	15.9 [11.8]	15.7 [12.9]	0.6895
Heart Rate	117.4 [40.4]	117.6 [39.8]	0.8381
Respiratory Rate	29.0 [13.9]	28.8 [13.9]	0.6250
Temperature (°C)	37.7 [1.8]	37.8 [1.8]	0.2379
$PAO_2/(0.01*FiO_2)$	215.5 [113.5]	215.8 [113.8]	0.9454
Albumin	3.1 [0.9]	3.0 [0.7]	0.3204
Hematocrit	31.0 [7.8]	31.1 [7.8]	0.6135
Bilirubin	2.3 [5.1]	2.1 [4.1]	0.1405
Creatinine	2.3 [2.2]	2.2 [2.1]	0.5349
Sodium	136.7 [7.9]	136.7 [7.9]	0.8792
Potassium	4.1 [1.1]	4.0 [1.0]	0.1329
$PACO_2$	37.1 [10.6]	37.6 [10.9]	0.1700
Serum Ph	7.4 [0.1]	7.4 [0.1]	0.9767
2nd Dx: Respiratory Diagnosis	35.2(%) [0.478]	38.0(%) [0.486]	0.0516
2nd Dx: Orthopedic Dx	0.2(%) [0.047]	0.0(%) [0.000]	0.0412
2nd Dx: Neurological Dx	8.6(%) [0.280]	9.7(%) [0.296]	0.1995
2nd Dx: GI Dx	16.0(%) [0.367]	14.7(%) [0.354]	0.2111
2nd Dx: Renal Dx	6.1(%) [0.240]	5.4(%) [0.227]	0.3439
2nd Dx: Metabolic Dx	5.2(%) [0.223]	4.8(%) [0.214]	0.5248
2nd Dx: Hematological Dx	7.0(%) [0.256]	7.4(%) [0.262]	0.6235
2nd Dx: Sepsis	21.5(%) [0.411]	21.0(%) [0.407]	0.6805
2nd Dx: Trauma	1.0(%) [0.101]	1.2(%) [0.108]	0.6611

Note: Each entry shows the mean and standard deviation (in brackets) for each variable.

Figure 1: % Catheterized by Day-of-Week of Admission by Diagnosis




Figure 2: Mortality Rates for Weekend vs. Weekday Admissions




Figure 3: OLS Treatment Effect




Figure 4: Bivariate Probit Treatment Effect




Figure 5: Manski (1990) and Shaikh and Vytlačil (2005) Bounds




Figure 6: PQD Bounds




Figure 7: PQD Bounds Without Instruments

